

### AT A GLANCE

# Infection Prevention and Control (IPAC) Recommendations for Monkeypox in Health Care Settings

Published: May 2022

### **Transmission**

Monkeypox virus can be transmitted from animals-to-humans (i.e., zoonotic transmission) or person-to-person by contact with infected lesions, skin scabs, body fluids or respiratory secretions. It can also be transmitted by contact with materials contaminated with the virus (e.g., clothing, bedding).<sup>1-4</sup>

Historically, there has been limited person-to-person transmission.<sup>3,4</sup> The primary mode of person-to-person transmission has been through respiratory secretions and direct contact with skin lesions or a patient's items that have been contaminated.<sup>2,4</sup> However, given the respiratory system involvement during infection, the possible transmission during the prodromal period and similarities to variola virus (smallpox), the potential for airborne transmission has been suggested.<sup>1-4</sup>

The incubation period averages 7 to 14 days (range 5 to 21 days). A person is most commonly contagious from the onset of initial lesions (typically on the tongue and in the mouth), until scabs have fallen off and new skin present. Some cases may be contagious during their early set of symptoms (prodrome) such as fever, malaise, headache before the rash develops.<sup>2</sup>

## IPAC Precautions in All Health Care Settings

In addition to Routine Practices, the following Additional Precautions - Airborne/Droplet/Contact Precautions are to be used.<sup>1</sup>

#### **Room Placement:**

- Airborne isolation rooms (AIR) with negative pressure ventilation.<sup>1,5</sup>
- When AIRs are not available, the patient can be placed in a single room with the door closed with dedicated toileting facility.
- If neither option is feasible, then precautions should be taken to minimize exposure to surrounding individuals such as having the patient don a medical mask over their nose and mouth as tolerated and covering exposed skin lesions with clothing, sheet or gown as best as possible.

Hand hygiene as per the Four Moments of Hand Hygiene.<sup>6</sup>

#### Personal Protective Equipment (PPE) for health care workers:

- Fit-tested and seal checked N-95 respirator (or equivalent); perform seal check after donning N95 respirator.
- Gloves
- Gown
- Eye protection (e.g., face shields or goggles)

**Duration:** Additional Precautions are maintained until all scabs have fallen off and new skin is present.

**Patient Transport:** Have the patient wear clean clothes/gown, wash their hands, wear a medical mask and cover their lesions to the best extent possible for transport.

**Laundry:** Special handling of linen from patients on Additional Precautions is not required. Care should be taken in the handling of soiled linen to prevent dispersal of microorganisms.<sup>1,5</sup>

**Waste disposal:** Containment and disposal of contaminated waste (e.g., dressings) in accordance with facility-specific guidelines for infectious waste.

**Environmental cleaning:** Healthcare-grade cleaning and disinfecting agents, with a Drug Identification Number (DIN) are appropriate for cleaning and disinfection of environmental surfaces and shared equipment in the patient care environment. Follow the manufacturer's recommendations for dilution and contact time.<sup>1,5</sup>

### References

- Public Health Agency of Canada. Routine practices and additional precautions for preventing
  the transmission of infection in healthcare settings [Internet]. Ottawa, ON: Her Majesty the
  Queen in Right of Canada; 2017 [modified 2017 Sep 26; cited 2022 May 19]. Available from:
  <a href="https://www.canada.ca/en/public-health/services/publications/diseases-conditions/routine-practices-precautions-healthcare-associated-infections.html">https://www.canada.ca/en/public-health/services/publications/diseases-conditions/routine-practices-precautions-healthcare-associated-infections.html</a>
- 2. Centers for Disease Control and Prevention. Monkeypox infection control: hospital [Internet]. Atlanta, GA: Centers for Disease Control and Prevention; 2015 [cited 2022 May 19]. Available from: https://www.cdc.gov/poxvirus/monkeypox/clinicians/infection-control-hospital.html
- 3. World Health Organization. Monkeypox [Internet]. Geneva: World Health Organization; 2022 [cited 2022 May 19]. Available from: <a href="https://www.who.int/news-room/fact-sheets/detail/monkeypox">https://www.who.int/news-room/fact-sheets/detail/monkeypox</a>
- 4. Public Health England. Monkeypox: information for primary care [Internet]. London: Crown Copyright; 2019 [cited 2022 May 19]. Available from:

  <a href="https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\_data/file/850059/Monkeypox\_information\_for\_primary\_care.pdf">https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\_data/file/850059/Monkeypox\_information\_for\_primary\_care.pdf</a>
- Ontario Agency for Health Protection and Promotion, Provincial Infectious Diseases Advisory Committee. Routine Practices and Additional Precautions in All Health Care Settings [Internet].
   3rd edition. Toronto, ON: Queen's Printer for Ontario; November 2012. [cited 2022 May 20]
   Available from: <a href="https://www.publichealthontario.ca/-/media/Documents/B/2012/bp-rpap-healthcare-settings.pdf?sc\_lang=en">https://www.publichealthontario.ca/-/media/Documents/B/2012/bp-rpap-healthcare-settings.pdf?sc\_lang=en</a>
- 6. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Hand hygiene [Internet]. Toronto, ON: Queen's Printer for Ontario; 2022 [cited 2022 May 19]. Available from: <a href="https://www.publichealthontario.ca/en/Health-Topics/Infection-Prevention-Control/Hand-Hygiene">https://www.publichealthontario.ca/en/Health-Topics/Infection-Prevention-Control/Hand-Hygiene</a>

### Citation

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Infection prevention and control (IPAC) recommendations for monkeypox in health care settings. Toronto, ON: Queen's Printer for Ontario; 2022

### Disclaimer

This document was developed by Public Health Ontario (PHO). PHO provides scientific and technical advice to Ontario's government, public health organizations and health care providers. PHO's work is guided by the current best available evidence at the time of publication. The application and use of this document is the responsibility of the user. PHO assumes no liability resulting from any such application or use. This document may be reproduced without permission for non-commercial purposes only and provided that appropriate credit is given to PHO. No changes and/or modifications may be made to this document without express written permission from PHO.

### **Public Health Ontario**

Public Health Ontario is an agency of the Government of Ontario dedicated to protecting and promoting the health of all Ontarians and reducing inequities in health. Public Health Ontario links public health practitioners, front-line health workers and researchers to the best scientific intelligence and knowledge from around the world.

For more information about PHO, visit <u>publichealthontario.ca</u>.

©Queen's Printer for Ontario, 2022

